

Dr. Richard Bartlett: Evidence Budesonide Is Safe, Effective Early and Late Treatment for COVID-19

- I. Dr. Bartlett's Published Protocol in Globaljournals.org
- II. U.K. STOIC Trial
- III. U.K. PRINCIPLE Trial
- IV. Brazil TOGETHER Trial
- V. Saudi Journal of Anesthesia Study
- VI. WHO Lists Budesonide as Essential Medicine
- VII. U.K. Department of Health Recommends Budesonide for Adults 50 or Over
- VIII. Australia Recommends Budesonide for Adults with COVID-19
- IX. India Ministry of Health and Family Welfare Recommends Budesonide for Adults with COVID-19
- X. Antiviral Effect of Budesonide Against COVID-19
- XI. FDA Rules Prohibit Agency from Granting Emergency Use Authorization (EUA) for Experimental COVID-19 Vaccines Because Budesonide Works

I. Dr. Bartlett's Published Protocol in Globaljournals.org

A 2020 [publication](#)¹ in *Global Journal of Science Frontier Research* titled "SARS-Cov-2 and the Case for Empirical Treatment" written by Doctors Richard Bartlett and Alexandria Watkins represented an "outpatient case study that examines two patients in the United States with unique cases that involve oncology and Severe Acute Respiratory Syndrome Coronavirus-2 (SARSCoV- 2), also known as COVID-19." The study "affirmed that an empirical treatment protocol with nebulized budesonide and the efficacy of treating symptomatic patients earlier rather than later has significant implications." The treatment plan even became "more effective by increasing the dosage and frequency of nebulized budesonide," according to the authors. A successful empirical treatment plan² included 0.5mg of nebulized budesonide "twice daily" and "clarithromycin (Biaxin) 500mg tab twice daily for ten days, Zinc 50mg tab twice daily, and aspirin 81mg tab daily." Bartlett's paper noted that there is a "decreased risk of pneumonia in COPD patients who use nebulized budesonide." "Secondary bacterial infection of the lung (pneumonia) was extremely common in patients with COVID-19," according to a report from Northwestern Medicine,³ citing data from an April 2023 study in *The Journal of Clinical Investigation*.⁴

II. U.K. STOIC Trial

A July 2021 [publication](#)⁵ in peer-reviewed medical journal *The Lancet Respiratory Medicine* "tested if inhaled glucocorticoids would be an effective treatment for early COVID-19." The study authors from Oxford University found that "[e]arly administration of inhaled budesonide reduced the likelihood of needing urgent medical care and reduced time to

¹ Richard P. Bartlett and Alexandria Watkins, "SARS-Cov-2 and the Case for Empirical Treatment," *Global Journal of Science Frontier Research*, July 2020, <https://journalofscience.org/index.php/GJSFR/article/view/2773>.

² Patients radically recovered: **EXHIBIT A:** Witness #1 Richard Jones and #2 Witness Brenda Jones; **EXHIBIT B:** Witness # 1 Daniel Cobarrubio (w/medical records) and #2 Anita Cobarrubio (American Faith coverage of Daniel's [story](#)).

³ Marla Paul, "Secondary Bacterial Pneumonia Drove Many COVID-19 Deaths," Northwestern Medicine, May 2023, <https://news.feinberg.northwestern.edu/2023/05/05/secondary-bacterial-pneumonia-drove-many-covid-19-deaths/#:~:text=By%20applying%20machine%20learning%20to%20medical%20record%20data%2C,results%20published%20in%20the%20Journal%20of%20Clinical%20Investigation>.

⁴ Catherine A. Gao *et al.*, "Machine learning links unresolving secondary pneumonia to mortality in patients with severe pneumonia, including COVID-19," *The Journal of Clinical Investigations*, April 2023, <https://www.jci.org/articles/view/170682>.

⁵ Sanjay Ramakrishnan *et al.*, "Inhaled budesonide in the treatment of early COVID-19 (STOIC): a phase 2, open-label, randomised controlled trial," *The Lancet Respiratory Medicine*, July 2021, [https://www.thelancet.com/journals/lanres/article/PIIS2213-2600\(21\)00160-0/fulltext#%20](https://www.thelancet.com/journals/lanres/article/PIIS2213-2600(21)00160-0/fulltext#%20).

recovery after early COVID-19.” The study represented an “open-label, parallel-group, phase 2, randomised controlled trial.” It concluded that “inhaled budesonide, when given to adults with early COVID-19, reduced the likelihood of requiring urgent care, emergency department consultation, or hospitalisation.” Budesonide administration allowed for “a quicker resolution of fever, a known poor prognostic marker in COVID-19.” The authors also revealed that “[s]elf-reported and questionnaire-reported symptom resolution was faster.” Moreover, there were “fewer participants with persistent COVID-19 symptoms at days 14 and 28 after budesonide therapy compared with usual care.” The author’s believe their study represents “the first interventional trial to study the efficacy of inhaled corticosteroids in early COVID-19 illness.” This STOIC trial “potentially provides the first easily accessible effective intervention in early COVID-19,” according to the researchers. “By assessing health-care resource use, the study provides an exciting option to help with the worldwide pressure on health-care systems due to the COVID-19 pandemic,” they write. “Data from this study also suggest a potentially effective treatment to prevent the long-term morbidity from persistent COVID-19 symptoms.”

A February 2021 University of Oxford [publication](#)⁶ titled “Common asthma treatment reduces need for hospitalisation in COVID-19 patients, study suggests” confirmed that the Lancet’s STOIC trial did find that early treatment with budesonide “appears to significantly reduce the need for urgent care and hospitalisation in people with COVID-19” and “reduced recovery time.” Inhaled budesonide “reduced the relative risk of requiring urgent care or hospitalisation by 90% in the 28-day study period,” Oxford states. Participants taking budesonide “also had a quicker resolution of fever, symptoms and fewer persistent symptoms after 28 days.” Oxford referenced Professor Mona Bafadhel of the University’s Nuffield Department of Medicine, who said she was “heartened that a relatively safe, widely available and well studied medicine such as an inhaled steroid could have an impact on the pressures we are experiencing during the pandemic.” Bafadhel, a Respiratory Consultant also working at the Oxford University Hospitals NHS Foundation Trust, commented on budesonide’s ability to reduce persistent COVID symptoms, calling it “an important finding.” “I am encouraged to see the reduction in persistent symptoms at 14 and 28 days after treatment with budesonide. Persistent symptoms after the initial COVID-19 illness have emerged as a long-term problem. Any intervention which could address this would be a major step forward,” she stated.

III. U.K. PRINCIPLE Trial

An August 2021 [publication](#)⁷ in *The Lancet* titled “Inhaled budesonide for COVID-19 in people at high risk of complications in the community in the UK (PRINCIPLE): a randomised, controlled, open-label, adaptive platform trial,” based on [Oxford University](#)’s⁸ ‘Platform Randomised Trial of Interventions against COVID-19 in Older People’ aimed to “establish whether inhaled budesonide reduces time to recovery and COVID-19-related hospital admissions or deaths among people at high risk of complications in the community.” This PRINCIPLE test represented a “multicentre, open-label, multi-arm, randomised, controlled, adaptive platform trial done remotely from a central trial site and at primary care centres in the UK,” according to the authors. Budesonide conferred “a benefit in time to first self-reported recovery” and lessened “hospital admission or death outcome.” The authors conclude, “Inhaled budesonide improves time to recovery, with a chance of also reducing hospital admissions or deaths ... in people with COVID-19 in the community who are at higher risk of complications.” While the authors noted that results pertaining to hospitalization and death “did not meet the superiority threshold,” they clarified that “this might have been due to the rapid decrease in rate of hospital admissions or deaths in March and April, 2021, in the UK, because of the vaccination programme and lockdown measures.” They went on to emphasize, “Overall, the consistency of these findings across both primary and secondary endpoints provides the strongest evidence thus far of an effective, safe, cheap, and readily available treatment for COVID-19 in the community.” This PRINCIPLE trial “is the largest randomised trial thus far to assess inhaled budesonide for community treatment of COVID-19.” It confirmed that “inhaled budesonide reduced COVID-19-related emergency assessments or hospital admissions, compared with usual care, and self-reported recovery favoured budesonide by 1 day.” The study authors also affirmed that “[s]everal randomised trials have shown that systemic corticosteroids reduce mortality among people admitted to hospital with COVID-19, with the RECOVERY trial finding greatest benefit in mechanically ventilated patients.” The PRINCIPLE

⁶ “Common asthma treatment reduces need for hospitalisation in COVID-19 patients, study suggests,” University of Oxford, February 2021, <https://www.ox.ac.uk/news/2021-02-09-common-asthma-treatment-reduces-need-hospitalisation-covid-19-patients-study>.

⁷ Ly-Mee Yu, “Inhaled budesonide for COVID-19 in people at high risk of complications in the community in the UK (PRINCIPLE): a randomised, controlled, open-label, adaptive platform trial,” *The Lancet*, August 2021, [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(21\)01744-X/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(21)01744-X/fulltext).

⁸ “Asthma drug budesonide shortens recovery time in non-hospitalised patients with COVID-19,” University of Oxford, April 2021, <https://www.ox.ac.uk/news/2021-04-12-asthma-drug-budesonide-shortens-recovery-time-non-hospitalised-patients-covid-19>.

trial “has provided evidence of a safe and cheap community treatment for COVID-19 that reduces symptom burden and enhances sustained recovery over 28 days, with a high probability of also reducing the need for hospital admission.” Inhaled budesonide “is available in many primary care settings and is included in the WHO list of essential medicines,” the authors highlighted (SEE VI). “Our study provides evidence that inhaled budesonide is an effective and safe treatment for people with COVID-19 in the community who are at increased risk of adverse outcomes.”

An August 2021 commentary [published](#)⁹ in *Annals of Medicine & Surgery* titled “Budesonide: A promising candidate therapeutic for early COVID-19” cited the above Lancet study, noting its “promising results of budesonide in treating patients with mild COVID-19 infection.” The authors underscored how “patients taking inhaled budesonide had a faster COVID-19 recovery time by 3 days than patients who received only usual care along with lower hospitalizations in the budesonide group than the usual care group.” They referenced the U.K. government’s conclusion that budesonide “can be considered (off label) on a case-by-case basis for symptomatic covid-19 positive patients aged 65 and over, or aged 50 or over with co-morbidities” (SEE VII). The authors state budesonide is a “a potent topical anti-inflammatory agent” in that it effectively translocates into the cell nucleus, prevents the expression of pro-inflammatory genes, increases the expression of anti-inflammatory genes, and “inhibits the eosinophil activation by increasing apoptosis and suppresses the activation of various inflammatory cells such as neutrophils, mast cells, macrophages, T-lymphocytes, and dendritic cells.” Budesonide administration therefore “leads to reduced airway inflammation and hyperreactivity resulting into inhibition of the bronchospasm and subsequently wheezing and coughing,” according to the authors. They conclude that inhaled budesonide “is a simple, safe, very well studied, widely available, and inexpensive corticosteroid which may prove crucial for mild COVID-19 cases.” Budesonide “could give healthcare workers more options in treating COVID-19 patients, especially as it is readily available in most of the primary healthcare settings and is listed as Essential Medicine in the World Health Organization’s List of Essential Medicines,” they conclude, and “can be used with ease even in comorbid, unwell, and potentially frail older patients.”

An April 2021 [publication](#)¹⁰ in *The British Medical Journal* (BMJ) titled “Covid-19: Budesonide shortens recovery time in patients not admitted to hospital, study finds” also recognized Oxford’s findings published in the Lancet, affirming that budesonide “can shorten the time it takes for people not admitted to hospital to recover from covid-19 by three days.” The BMJ piece emphasized how those treated with budesonide “also reported greater wellbeing after two weeks.” Less of those treated with budesonide “were admitted to hospital” than those who did not. The author for BMJ cited Fiona Watt, Executive Chair of the Medical Research Council, who said that “researchers involved in the Principle trial have overcome considerable logistical hurdles to set up a world leading rigorous drug trial in people’s homes.” Watt added, “We are now rewarded with the first inexpensive and widely available drug that can shorten recovery times for covid-19 patients in the community.” The author also cited Joint Chief Investigator Chris Butler, a south Wales GP and Professor of Primary Care from the University of Oxford, who said, “We therefore anticipate that medical practitioners around the world caring for people with covid-19 in the community may wish to consider this evidence when making treatment decisions.”

The Lancet PRINCIPLE study authors [published](#)¹¹ “Inhaled Budesonide for COVID-19 in People at Higher Risk of Complications in the Community: The UK National Community Randomi” in *Annals of Family Medicine* in January 2023. In that publication, they confirm that inhaled budesonide (800µg twice daily for 14 days) led to “shorter” time to first self-reported recovery compared to usual care and less hospitalizations and deaths compared to usual care.

IV. Brazil TOGETHER Trial

A May 2023 [publication](#)¹² in *Annals of Internal Medicine* titled “Oral Fluvoxamine With Inhaled Budesonide for Treatment of Early-Onset COVID-19” set out to “determine whether the combination of fluvoxamine and inhaled budesonide would increase treatment effects in a highly vaccinated population” in a randomized, placebo-controlled, adaptive platform trial. “Patients were randomly assigned to either fluvoxamine (100 mg twice daily for 10 days) plus inhaled budesonide (800

⁹ Shailesh Kumar Patel *et al.*, “Budesonide: A promising candidate therapeutic for early COVID-19,” *Annals of Medicine & Surgery*, August 2021, https://journals.lww.com/annals-of-medicine-and-surgery/Fulltext/2021/08000/Budesonide_A_promising_candidate_therapeutic_for.58.aspx.

¹⁰ Elisabeth Mahase, “Covid-19: Budesonide shortens recovery time in patients not admitted to hospital, study finds,” *The British Medical Journal* (BMJ), April 2021, <https://www.bmj.com/content/373/bmj.n957>.

¹¹ Richard Hobbs *et al.*, “Inhaled Budesonide for COVID-19 in People at Higher Risk of Complications in the Community: The UK National Community Randomi,” *Annals of Family Medicine*, January 2023, https://www.annfammed.org/content/21/Supplement_1/3859.

¹² Gilmar Reis *et al.*, “Oral Fluvoxamine With Inhaled Budesonide for Treatment of Early-Onset COVID-19,” *Annals of Internal Medicine*, May 2023, https://www.acpjournals.org/doi/full/10.7326/M22-3305?rfr_dat=cr_pub++0pubmed&url_ver=Z39.88-2003&rfr_id=ori%3Arid%3Acrossref.org.

mcg twice daily for 10 days) or matching placebos,” the study reads. The number of patients “observed in an emergency setting for COVID-19 for more than 6 hours or hospitalized due to COVID-19 was lower in the treatment group than the placebo group.” The authors conclude that treatment with oral fluvoxamine plus inhaled budesonide “among high-risk outpatients with early COVID-19 reduced the incidence of severe disease requiring advanced care.” The study was “among the first to evaluate a drug combination for treatment of ambulatory patients with COVID-19 in a randomized trial,” according to the authors. They “found a reduction in the composite end point for COVID-19 disease progression with a combination of oral fluvoxamine, 100 mg twice daily, and inhaled budesonide.” Budesonide and fluvoxamine “reduced the need for advanced medical care.” Moreover, the “number of serious adverse events associated with this combination therapy was lower than in the placebo group.” The authors confirm that “In conclusion, administration of the combination of fluvoxamine, 100 mg twice daily, and inhaled budesonide reduced the rate of COVID-19 progression resulting in prolonged observation in an emergency setting or hospitalization among outpatients with high risk for serious disease.”

V. Saudi Journal of Anesthesia Study

A January 2017 [publication](#)¹³ in *Saudi Journal of Anesthesia* “tested the hypothesis that nebulized budesonide would improve lung mechanics and oxygenation in patients with early acute lung injury (ALI) and/or acute respiratory distress syndrome (ARDS) during protective mechanical ventilation strategy without adversely affecting systemic hemodynamics.” Patients received “1 mg–2 ml budesonide suspension ... nebulized through the endotracheal tube” every “12 h for three successive days alongside with constant ventilator settings.” The study authors found that nebulized budesonide “improved oxygenation, peak, and plateau airway pressures and significantly reduced inflammatory markers (TNF- α , IL-1 β and IL-6) without affecting hemodynamics.” Budesonide “inhibits a variety of inflammatory cells, reduces the production of inflammatory mediators and consequently has a significant anti-inflammatory effect.” It “induces vasoconstriction, inhibits mucosal edema, reduces cell exudation, and prevents airway remodeling.” It can “obtund the sequelae of the acute phase response and reduces alveolar inflammation.” Budesonide “inhibits airway inflammation, alleviates edema, inhibits remodeling and promotes suctioning which maintains pulmonary function during mechanical ventilation.” It “reduces the adverse events associated with systemic corticosteroid administration such as increased blood glucose level, hypothalamic-pituitary-adrenal axis suppression, bone demineralization, perforated peptic ulcer, and altered immunity.” Budesonide “can act as a pulmonary protective agent during mechanical ventilation for ALI/ARDS patients.” Late COVID results in ARDS, this study proving that budesonide is an effective and safe treatment for ARDS.

VI. WHO Lists Budesonide as Essential Medicine

A July 2019 World Health Organization (WHO) [publication](#)¹⁴ titled “WHO model list of essential medicines - 21st list, 2019” recommends budesonide inhaled (aerosol) at “100 micrograms per dose” and “200 micrograms per dose” as one of its “medicines acting on the respiratory tract.”

VII. U.K. Department of Health Recommends Budesonide for Adults 50 or Over

An April 2021 U.K. Department of Health [alert](#)¹⁵ from Chief Medical Officer Dr. Michael McBride titled “COVID-19 THERAPEUTIC ALERT - INHALED BUDESONIDE FOR ADULTS (50 YEARS AND OVER) WITH COVID-19” stated that inhaled budesonide “can be considered (off-label) on a case-by-case basis for symptomatic COVID-19 positive patients aged 65 and over, or aged 50 or over with co-morbidities, in line with the published Interim Position Statement.”

The aforementioned April 2021 BMJ publication (SEE III) recognized the U.K. government health alert affirming that budesonide “can be considered” to treat COVID-19.

¹³ Hatem Saber Mohamed and Mona Mohamed Abdel Meguid, “Effect of nebulized budesonide on respiratory mechanics and oxygenation in acute lung injury/acute respiratory distress syndrome,” *Saudi Journal of Anesthesia*, January–March 2017, https://journals.lww.com/sjan/Fulltext/2017/11010/Effect_of_nebulized_budesonide_on_respiratory.3.aspx.

¹⁴ “WHO model list of essential medicines - 21st list, 2019,” World Health Organization, July 2019, <https://www.who.int/publications/i/item/WHOMVPEMPIAU2019.06>.

¹⁵ Michael McBride, U.K. Chief Medical Officer, “HSS(MD) 29/2021 – COVID-19 THERAPEUTIC ALERT – INHALED BUDESONIDE FOR ADULTS (50 YEARS AND OVER) WITH COVID-19,” U.K. Health and Social Care, April 2021, <https://gpmi.co.uk/hsc-updates/hssmd-29-2021-covid-19-therapeutic-alert-inhaled-budesonide-for-adults-50-years-and-over-with-covid-19/>.

VIII. Australia Recommends Budesonide for Adults with COVID-19

Australia's Department of Health, through its National Clinical Evidence Taskforce, [published](#)¹⁶ a document titled "COVID-19 medications for at risk people who do not require oxygen" in June 2022 that recommends budesonide at "800 µg inhaled twice daily for up to 14 days." Budesonide "can be considered as either a standalone therapy or as an additional therapy in patients already prescribed another early therapy," according to the document.

IX. India Ministry of Health and Family Welfare Recommends Budesonide for Adults with COVID-19

India's Ministry of Health and Family Welfare [published](#)¹⁷ a document titled "Clinical Management Protocol for COVID-19" in May 2021 that recommends budesonide "given via inhalers with spacer at a dose of 800 mcg twice daily for 5 to 7 days."

X. Antiviral Effect of Budesonide Against COVID-19

A July 2021 [publication](#)¹⁸ in *Viruses* titled "Antiviral Effect of Budesonide against SARS-CoV-2" analyzed the "potential antiviral activity of budesonide" and potentially indicates a "multimodal mode of action of budesonide against SARS-CoV-2 and COVID-19." The study results "suggest that treatment with budesonide reduces titers of SARS-CoV-2 and VOCs significantly while cell viability remains unaffected," according to the authors. They "observed significant reduction of viral titers for all viral variants in vitro when cells were treated with 25 µM budesonide."

XI. FDA Rules Prohibit Agency from Granting Emergency Use Authorization (EUA) for Experimental COVID-19 Vaccines Because Budesonide Works

The U.S. Food & Drug Administration (FDA) affirms that it "can use its Emergency Use Authorization (EUA) authority under section 564 of the Federal Food, Drug, and Cosmetic Act (FD&C Act) to allow the use of unapproved medical products, or unapproved uses of approved medical products, to diagnose, treat, or prevent serious or life-threatening diseases when certain criteria are met," according to the FDA website.¹⁹ This is "including that there are no adequate, approved, and available alternatives." Section 564 states that an EUA cannot be granted if "there is no adequate, approved, and available alternative to the product for diagnosing, preventing, or treating such disease or condition."²⁰ Budesonide is an "adequate, approved,²¹ and available alternative" to COVID-19 vaccines, thus meeting the FDA's section 564 criteria prohibiting the EUA of COVID vaccines. Therefore, experimental COVID vaccines EUAs are void and illegal. Present and future production of COVID vaccines and/or gene therapies must cease and desist. The FDA is currently egregiously violating its own rules.

¹⁶ "COVID-19 medications for at risk people who do not require oxygen," Australia's Department of Health through its National Clinical Evidence Taskforce, June 2022, <https://www.health.vic.gov.au/publications/covid-19-medications-for-at-risk-people-who-do-not-require-oxygen>.

¹⁷ "Clinical Management Protocol for COVID-19," Government of India Ministry of Health and Family Welfare, May 2021, <https://www.mohfw.gov.in/pdf/UpdatedDetailedClinicalManagementProtocolforCOVID19adultsdated24052021.pdf>.

¹⁸ Natalie Heinen *et al.*, "Antiviral Effect of Budesonide against SARS-CoV-2," *Viruses*, July 2021, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8310374/>.

¹⁹ "FAQs on Emergency Use Authorizations (EUAs) for Medical Devices Related to COVID-19," U.S. Food & Drug Administration, March 2023, <https://www.fda.gov/medical-devices/covid-19-emergency-use-authorizations-medical-devices/faqs-emergency-use-authorizations-euas-medical-devices-related-covid-19>.

²⁰ 21 U.S.C., United States Code, 2011 Edition, Title 21 - FOOD AND DRUGS, CHAPTER 9 - FEDERAL FOOD, DRUG, AND COSMETIC ACT, U.S. Government Publishing Office, <https://www.govinfo.gov/content/pkg/USCODE-2011-title21/html/USCODE-2011-title21-chap9.htm>.

²¹ Drug Approval Package, Budesonide inhalation powder, Company: AstraZeneca Pharmaceuticals, NDA: 021949, Approval Date: 7/12/2006, U.S. Food & Drug Administration (FDA), November 2008, https://www.accessdata.fda.gov/drugsatfda_docs/nda/2006/021949_budesonide_toc.cfm.

Bibliography

- 21 U.S.C. United States Code. 2011 Edition. Title 21 - FOOD AND DRUGS, CHAPTER 9 – FEDERAL FOOD, DRUG, AND COSMETIC ACT. U.S. Government Publishing Office. <https://www.govinfo.gov/content/pkg/USCODE-2011-title21/html/USCODE-2011-title21-chap9.htm>.
- “Asthma drug budesonide shortens recovery time in non-hospitalised patients with COVID-19.” University of Oxford. April 2021. <https://www.ox.ac.uk/news/2021-04-12-asthma-drug-budesonide-shortens-recovery-time-non-hospitalised-patients-covid-19>.
- Bartlett, Richard P. and Alexandria Watkins. “SARS-Cov-2 and the Case for Empirical Treatment.” *Global Journal of Science Frontier Research*. July 2020. <https://journalofscience.org/index.php/GJSFR/article/view/2773>.
- “Clinical Management Protocol for COVID-19.” Government of India Ministry of Health and Family Welfare. May 2021. <https://www.mohfw.gov.in/pdf/UpdatedDetailedClinicalManagementProtocolforCOVID19adultsdated24052021.pdf>.
- “Common asthma treatment reduces need for hospitalisation in COVID-19 patients, study suggests.” University of Oxford. February 2021. <https://www.ox.ac.uk/news/2021-02-09-common-asthma-treatment-reduces-need-hospitalisation-covid-19-patients-study>.
- “COVID-19 medications for at risk people who do not require oxygen.” Australia’s Department of Health through its National Clinical Evidence Taskforce. June 2022. <https://www.health.vic.gov.au/publications/covid-19-medications-for-at-risk-people-who-do-not-require-oxygen>.
- Drug Approval Package. Budesonide inhalation powder. Company: AstraZeneca Pharmaceuticals. NDA: 021949. Approval Date: 7/12/2006. U.S. Food & Drug Administration (FDA). November 2008. https://www.accessdata.fda.gov/drugsatfda_docs/nda/2006/021949_budesonide_toc.cfm.
- “FAQs on Emergency Use Authorizations (EUAs) for Medical Devices Related to COVID-19.” U.S. Food & Drug Administration. March 2023. <https://www.fda.gov/medical-devices/covid-19-emergency-use-authorizations-medical-devices/faqs-emergency-use-authorizations-euas-medical-devices-related-covid-19>.
- Gao, Catherine A. *et al.* “Machine learning links unresolving secondary pneumonia to mortality in patients with severe pneumonia, including COVID-19.” *The Journal of Clinical Investigations*. April 2023. <https://www.jci.org/articles/view/170682>.
- Heinen, Natalie *et al.* “Antiviral Effect of Budesonide against SARS-CoV-2.” *Viruses*. July 2021. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8310374/>.

- Hobbs, Richard *et al.*, “Inhaled Budesonide for COVID-19 in People at Higher Risk of Complications in the Community: The UK National Community Randomi.” *Annals of Family Medicine*. January 2023. https://www.annfammed.org/content/21/Supplement_1/3859.
- Mahase, Elisabeth. “Covid-19: Budesonide shortens recovery time in patients not admitted to hospital. study finds.” *The British Medical Journal* (BMJ). April 2021. <https://www.bmj.com/content/373/bmj.n957>.
- McBride, Michael. U.K. Chief Medical Officer. “HSS(MD) 29/2021 – COVID-19 THERAPEUTIC ALERT – INHALED BUDESONIDE FOR ADULTS (50 YEARS AND OVER) WITH COVID-19.” U.K. Health and Social Care. April 2021. <https://gpni.co.uk/hsc-updates/hssmd-29-2021-covid-19-therapeutic-alert-inhaled-budesonide-for-adults-50-years-and-over-with-covid-19/>.
- Mohamed, Hatem Saber and Mona Mohamed Abdel Meguid. “Effect of nebulized budesonide on respiratory mechanics and oxygenation in acute lung injury/acute respiratory distress syndrome.” *Saudi Journal of Anesthesia*. January–March 2017. https://journals.lww.com/sjan/Fulltext/2017/11010/Effect_of_nebulized_budesonide_on_respiratory.3.aspx.
- Patel, Shailesh Kumar *et al.* “Budesonide: A promising candidate therapeutic for early COVID-19.” *Annals of Medicine & Surgery*. August 2021. https://journals.lww.com/annals-of-medicine-and-surgery/Fulltext/2021/08000/Budesonide_A_promising_candidate_therapeutic_for.58.aspx.
- Paul, Marla. “Secondary Bacterial Pneumonia Drove Many COVID-19 Deaths.” *Northwestern Medicine*. May 2023. <https://news.feinberg.northwestern.edu/2023/05/05/secondary-bacterial-pneumonia-drove-many-covid-19-deaths/#:~:text=By%20applying%20machine%20learning%20to%20medical%20record%20data%2C,results%20published%20in%20the%20Journal%20of%20Clinical%20Investigation>.
- Ramakrishnan, Sanjay *et al.* “Inhaled budesonide in the treatment of early COVID-19 (STOIC): a phase 2, open-label, randomised controlled trial.” *The Lancet Respiratory Medicine*. July 2021. [https://www.thelancet.com/journals/lanres/article/PIIS2213-2600\(21\)00160-0/fulltext#%20](https://www.thelancet.com/journals/lanres/article/PIIS2213-2600(21)00160-0/fulltext#%20).
- Reis, Gilmar *et al.* “Oral Fluvoxamine With Inhaled Budesonide for Treatment of Early-Onset COVID-19.” *Annals of Internal Medicine*. May 2023. https://www.acpjournals.org/doi/full/10.7326/M22-3305?rfr_dat=cr_pub++0pubmed&url_ver=Z39.88-2003&rfr_id=ori%3Arid%3Acrossref.org.
- “WHO model list of essential medicines - 21st list, 2019.” World Health Organization. July 2019. <https://www.who.int/publications/i/item/WHOMVPEMPIAU2019.06>.

Yu, Ly-Mee *et al.* “Inhaled budesonide for COVID-19 in people at high risk of complications in the community in the UK (PRINCIPLE): a randomised, controlled, open-label, adaptive platform trial.” *The Lancet*. August 2021. [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(21\)01744-X/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(21)01744-X/fulltext).